






Effects of high intensity non-ionizing terahertz radiation on human skin fibroblasts: supplement

DMITRY S. SITNIKOV,^{1,*}  INNA V. ILINA,¹  VERONIKA A. REVKOVA,² SERGEY A. RODIONOV,³ SVETLANA A. GUROVA,⁴ RIMMA O. SHATALOVA,⁴ ALEXEY V. KOVALEV,³  ANDREY V. OVCHINNIKOV,¹  OLEG V. CHEFONOV,¹  MIKHAIL A. KONOPLYANNIKOV,^{2,5} VLADIMIR A. KALSIN,² AND VLADIMIR P. BAKLAUSHEV²

¹Joint Institute for High Temperatures of the Russian Academy of Sciences, Moscow, Russia

²Federal Research and Clinical Center of Specialized Medical Care and Medical Technologies FMBA of Russia, Moscow, Russia

³N. N. Priorov National Medical Research Center of Traumatology and Orthopedics, Moscow, Russia

⁴National Research nuclear University MEPhI Obninsk Institute for Nuclear Power Engineering, Obninsk, Russia

⁵Institute for Regenerative Medicine, Sechenov First Moscow State Medical University, Moscow, Russia

*Sitnik.ds@gmail.com

This supplement published with Optica Publishing Group on 26 October 2021 by The Authors under the terms of the [Creative Commons Attribution 4.0 License](https://creativecommons.org/licenses/by/4.0/) in the format provided by the authors and unedited. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI.

Supplement DOI: <https://doi.org/10.6084/m9.figshare.16837666>

Parent Article DOI: <https://doi.org/10.1364/BOE.440460>

Effects of high intensity non-ionizing terahertz radiation on human skin fibroblasts: supplemental document

In the course of this work, we also studied the activity of heat shock proteins as a result of thermal exposure to various temperatures. It is noteworthy that the activation of HSP70 took place in both hypothermia and hyperthermia (Fig. S1). However, during hyperthermia, the expression of heat shock proteins increased significantly. A good fluorescence signal was observed starting from 40°C, it increased maximally at 42.5°C and did not change at 45°C. However, at 45°C, cell deformation and the appearance of some parts of apoptotic cells in culture were observed.




Fig. S1. Expression of heat shock proteins (HSP70) in human skin fibroblasts at different temperatures. (a) 4°C, (b) 40°C, (c) 42.5°C, (d) 45°C.